






# ESOPHAGUS CANCER BY RACE IN NORTHEAST OHIO, 2016 - 2020

September 2023

## Age-Adjusted Incidence Rate



	All	NH-White	NH-Black	Hispanic	Other**
<b>Catchment Area</b>	5.2 (4.9 – 5.5)	5.6 (5.3 – 6.0)	3.3 (2.7 – 4.0)	N/A***	2.9 (1.5 – 5.4)
<b>Ohio</b>	5.6 (5.5 – 5.8)	6.0 (5.8 – 6.2)	3.6 (3.2 – 4.1)	1.5 (0.9 – 2.5)	3.0 (2.1 – 4.2)
<b>National</b>	4.5 (4.5 – 4.5)	5.0 (5.0 – 5.1)	3.5 (3.4 – 4.6)	2.7 (2.6 – 2.7)	2.4 (2.3 – 2.5)

**Direct Age Adjusted Incidence Rate (95% Confidence Interval)**

Average annual rate per 100,000, age adjusted to the 2000 US standard population.

\* Ashland, Ashtabula, Cuyahoga, Erie, Geauga, Huron, Lake, Lorain, Mahoning, Medina, Portage, Stark, Summit, Trumbull, and Wayne

\*\* American Indian or Alaskan Native, and Asian or Pacific Islander

\*\*\* N/A-indicates where rates may be unstable for case counts less than eleven

Ohio Cancer Incidence Surveillance System, Ohio Department of Health. Official End of Year from 1996 – 2020

National Program of Cancer Registries and Surveillance, Epidemiology and End Results Program SEER\*Stat Database: NPCR and SEER Incidence - U.S. Cancer Statistics Public Use Research Database, 2022 Submission (2001-2020). United States Department of Health and Human Services, Centers for Disease Control and Prevention and National Cancer Institute. Released June 2023. Accessed at [www.cdc.gov/cancer/uscs/public-use](http://www.cdc.gov/cancer/uscs/public-use).