






ESOPHAGUS CANCER BY RACE IN NORTHEAST OHIO, 2016 - 2020

September 2023

Age-Adjusted Mortality Rate



	All	NH-White	NH-Black	Hispanic	Other**
Catchment Area	4.6 (4.3 – 4.8)	4.9 (4.6 – 5.2)	2.7 (2.2 – 3.4)	3.2 (1.8 – 5.1)	N/A
Ohio	4.9 (4.7 – 5.0)	5.2 (5.0 – 5.3)	3.1 (2.7 – 3.5)	2.8 (1.9 – 3.9)	2.4 (1.6 – 3.5)
National	3.8 (3.8 – 3.8)	4.3 (4.3 – 4.4)	2.9 (2.8 – 3.0)	1.9 (1.9 – 2.0)	1.7 (1.6 – 1.8)

Direct Age Adjusted Mortality Rate (95% Confidence Interval)

Average annual rate per 100,000, age adjusted to the 2000 US standard population.

* Ashland, Ashtabula, Cuyahoga, Erie, Geauga, Huron, Lake, Lorain, Mahoning, Medina, Portage, Stark, Summit, Trumbull, and Wayne

** American Indian or Alaskan Native, and Asian or Pacific Islander

*** N/A-indicates where rates may be unstable for case counts less than eleven

Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1990-2020) < Katrina/Rita Population Adjustment > - Linked To County Attributes - Total U.S., 1969-2020 Counties, National Cancer Institute, DCCPS, Surveillance Research Program, released June 2022. Underlying mortality data provided by NCHS (www.cdc.gov/nchs).