


Even though African Americans comprise 17% of the population of both races, they comprise 50% of excess incident cases (Non-Hispanic Black vs Non-Hispanic White: 62 vs 61), and 30% of excess deaths (Non-Hispanic Black vs Non-Hispanic White: 101 vs 237).

Comparison of the Catchment Area and U.S. Average Annual Age-adjusted Incidence Rates by Cancer Site and Race, 2016-2020



Notes:

1. Cancers are included if the rate is among the top 20, and the absolute rate difference between catchment area and U.S. is at least 5%.
2. Catchment Area include 15 counties in Northeast Ohio: Ashland, Ashtabula, Cuyahoga, Erie, Geauga, Huron, Lake, Lorain, Mahoning, Medina, Portage, Stark, Summit, Trumbull, and Wayne Counties.
3. Rates for both the catchment area and the U.S. were specific to the respective racial/ethnic group.
4. Excessive cases (per year) is estimated by multiply the population of the catchment area by the difference in incidence rates between the catchment area and the US.

* Rates for uterus, cervix, and ovary cancers are limited to females.

** IBD: Intrahepatic Bile Duct.

Sources:

1. Catchment area data: Ohio Public Health Information Warehouse, Ohio Department of Health. Cancer Incidence Data (1996-2020). Access at <https://publicapps.odh.ohio.gov/EDW/DataBrowser/Browse/StateLayoutLockdownCancers>
2. US data: National Program of Cancer Registries and Surveillance, Epidemiology and End Results Program SEER*Stat Database: NPCR and SEER Incidence - U.S. Cancer Statistics Public Use Research Database, 2022 Submission (2001-2020). United States Department of Health and Human Services, Centers for Disease Control and Prevention and National Cancer Institute. Released June 2023. Accessed at www.cdc.gov/cancer/uscs/public-use