




LUNG AND BRONCHUS CANCER BY RACE IN NORTHEAST OHIO, 2016 - 2020

September 2023

Age-Adjusted Incidence Rate



	All	NH-White	NH-Black	Hispanic	Other**
Catchment Area	61.4 (60.5 – 62.3)	61.6 (60.6 – 62.6)	68.7 (65.9 – 71.6)	25.1 (20.9 – 30.1)	40.4 (34.4 – 47.3)
Ohio	64.7 (64.2 – 65.3)	65.7 (65.1 – 66.3)	68.0 (66.2 – 70.0)	22.2 (19.4 – 25.2)	38.2 (34.7 – 41.9)
National	54.2 (54.1 – 54.3)	58.7 (58.6 – 58.9)	55.7 (55.4 – 56.0)	27.6 (27.4 – 27.9)	35.0 (34.7 – 35.4)

Direct Age Adjusted Incidence Rate (95% Confidence Interval)

Average annual rate per 100,000, age adjusted to the 2000 US standard population.

* Ashland, Ashtabula, Cuyahoga, Erie, Geauga, Huron, Lake, Lorain, Mahoning, Medina, Portage, Stark, Summit, Trumbull, and Wayne

** American Indian or Alaskan Native, and Asian or Pacific Islander

Ohio Cancer Incidence Surveillance System, Ohio Department of Health. Official End of Year from 1996–2020

National Program of Cancer Registries and Surveillance, Epidemiology and End Results Program SEER*Stat Database: NPCR and SEER Incidence - U.S. Cancer Statistics Public Use Research Database, 2022 Submission (2001-2020). United States Department of Health and Human Services, Centers for Disease Control and Prevention and National Cancer Institute. Released June 2023. Accessed at www.cdc.gov/cancer/uscs/public-use.