






LUNG AND BRONCHUS CANCER BY RACE IN NORTHEAST OHIO, 2016 - 2020

September 2023

Age-Adjusted Mortality Rate



	All	NH-White	NH-Black	Hispanic	Other**
Catchment Area	40.7 (39.9 – 41.5)	40.8 (39.9 – 41.6)	46.3 (44.0 – 48.7)	16.4 (13.1 – 20.3)	23.7 (19.0 – 29.2)
Ohio	42.9 (42.5 – 43.4)	43.7 (43.2 – 44.2)	44.8 (43.2 – 46.3)	15.6 (13.3 – 18.1)	20.4 (17.8 – 23.3)
National	35.0 (34.9 – 35.0)	38.0 (37.9 – 38.1)	37.2 (37.0 – 37.5)	15.4 (15.2 – 15.6)	21.0 (20.8 – 21.3)

Direct Age Adjusted Mortality Rate (95% Confidence Interval)

Average annual rate per 100,000, age adjusted to the 2000 US standard population.

* Ashland, Ashtabula, Cuyahoga, Erie, Geauga, Huron, Lake, Lorain, Mahoning, Medina, Portage, Stark, Summit, Trumbull, and Wayne

** American Indian or Alaskan Native, and Asian or Pacific Islander

Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1990-2020) < Katrina/Rita Population Adjustment > - Linked To County Attributes - Total U.S., 1969-2020 Counties, National Cancer Institute, DCCPS, Surveillance Research Program, released June 2022. Underlying mortality data provided by NCHS (www.cdc.gov/nchs).